arXiv:1207.5194v2 [hep-ph] 31 Aug 2012

The $O(4)$ violating terms of the multi-Higgs-doublet model potential cannot be excluded by imposing a symmetry on the Lagrangian

M. Aa. Solberg[1, 2]

[1]*Program for Mechanical and Logistic Engineering, HiST, N–7004 Trondheim, Norway*[*]
[2]*Department of Physics, NTNU, N–7491 Trondheim, Norway*

We prove there is no symmetry which, imposed on the general charge conjugation (C) invariant multi-Higgs-doublet model (NHDM), excludes exactly the $O(4)$ violating terms from the potential. This is in part done by determining the exact subsymmetries of the custodial $SO(4)$ symmetry of the standard model (SM) scalar Lagrangian. Furthermore, the Higgs family symmetries of the quadratic part of the general C invariant NHDM potential will also be symmetries of the $O(4)$ violating terms of the potential, and hence cannot be applied to remove the $O(4)$ violating terms from the potential. Finally, negative determinant elements of $O(4)$ cannot be applied to expel the $O(4)$ breaking terms from the NHDM potential either: We show there is no symmetries of the kinetic Higgs terms in the negative determinant part of $O(4)$. This is also the case in the limit $g' \to 0$. Hence the custodial $SO(4)$ symmetry is the largest symmetry group of the scalar sector of the SM in the limit $g' \to 0$.

I. INTRODUCTION

A recent analysis of data from the LHC has proven the existence of a Higgs-like boson with mass at about 126 GeV [1, 2], while future investigations might give an indication of a sector of scalar particles beyond the single Higgs boson postulated by the standard model (SM). A natural extension of the SM involving several scalar particles is the SM augmented by several Higgs doublets, resulting in what is denoted multi-Higgs-doublet models (NHDM). In these models, some of the Higgs bosons should be responsible for the generation of the masses of fermions and the electroweak bosons [3–6]. Other Higgs particles might incorporate the dark matter [7–14]. In addition, the NHDM naturally accommodates CP violation [13, 15–19].

What is usually referred to as the custodial symmetry is an approximate $SO(3)$ symmetry of the SM: The symmetry is exact in the limit $g' \to 0$ (where g' is the hypercharge coupling constant), in which the gauge bosons $W^{\pm}, Z$ form a triplet (with identical masses). The name "custodial" was chosen since the symmetry guards the tree-level relation [20]

$$\rho \equiv \frac{m_W^2}{m_Z^2 \cos\theta_W} = 1, \tag{1}$$

from radiative corrections. The radiative corrections are proportional to g'^2, and hence vanish in the custodial symmetric limit $g' \to 0$ [21]. What we refer to as "the custodial $SO(4)$ symmetry" is the $(SU(2)_L \times SU(2)_R)/\mathbb{Z}_2 \cong SO(4)$ symmetry which contains the custodial $SO(3)$ symmetry, see sec. II. The custodial $SO(4)$ symmetry is, before spontaneous symmetry breaking, exact in the limit $g' \to 0$, for the scalar sector of the SM.

In an earlier paper [22] we investigated symmetry properties of the NHDM. Here we saw that the most general C invariant NHDM potential has an $O(4)$ symmetry (extending the custodial $SO(4)$ symmetry) only broken by certain quartic terms in the Higgs fields, of the type $\widehat{C}^2$, cf. eq. (7) below. The terms $\widehat{C}$ are odd under charge conjugation C, and hence terms quadratic in $\widehat{C}$ are invariant under C. We found that, in case of real vacuum expectation values (VEVs), the presence of terms $\lambda^{(3)}\widehat{C}^2$, cf. (9), violates a symmetry between the charged and C odd Higgs sectors. We argued that the C odd and charged sectors will get identical mass spectra, in case the parameters of the type $\lambda^{(3)}$ initially are set to zero. But when it comes to scattering processes, terms of the type $\widehat{C}^2$ may show up as counterterms during renormalization, even though their corresponding parameters $\lambda^{(3)}$ initially are set to zero [22].

Thus a question arises: Is it possible to find a discrete symmetry D which is a symmetry of the C-invariant Higgs sector, except that it expels the $O(4)$-violating terms, i.e. the terms of the type $\widehat{C}^2$, from the NHDM potential when it is imposed on the NHDM Lagrangian? If there exists such a symmetry, we may impose this symmetry on the NHDM Lagrangian, and hence avoid terms of the type $\widehat{C}^2$ both in the original Lagrangian and as counterterms under the renormalization procedure.

---

[*]Electronic address: marius.solberg@hist.no

The discrete symmetry D has evidently to lie beyond the symmetry group of $\widehat{C}^2$ (which we denoted P), but has to be a symmetry of the (other) terms of the Higgs Lagrangian. Hence it has to be an element of $O(4)$, which is the greatest possible symmetry group of a Lagrangian which can be expressed by real fields organized in quadruplets, provided that the NHDM potential is complicated enough so that different Higgs quadruplets cannot be transformed into each other. As we considered in [22], the symmetry group of the kinetic Higgs terms may be larger than $O(4)$ if fields with different Higgs indices can be transformed into each other: We here showed that the kinetic terms where invariant under $SU(2) \times U(N)$ in the case of N complex Higgs doublets. The $U(N)$ component will generally not be a symmetry of the Higgs potential, since the Higgs fields generally will occur in an asymmetrical manner in a potential. An element of this $U(N)$ component of the symmetry group of the kinetic terms, that also is a symmetry of a specific NHDM potential, is denoted a Higgs family (HF) symmetry of that potential [23]. In appendix A we show the only HF symmetry of the quadratic (in the Higgs fields) part of the general C invariant NHDM potential is an $U(1)$ transformation. This is a symmetry of all terms in a NHDM potential, hence a HF symmetry cannot be imposed to expel the terms of the type $\widehat{C}^2$ (and these terms only) from the most general C invariant NHDM potential.

Hence we can state that a symmetry transformation of a NHDM Lagrangian with a sufficiently complicated potential has to be an element of $O(4)$ containing the custodial $SO(4)$ symmetry. In section V B 1 we show that no element of $O^-(4)$ (orthogonal matrices with negative determinant) is a symmetry transformation of the kinetic Higgs terms, and hence an exact (discrete or continuous) symmetry of the scalar SM Lagrangian has to be an exact subsymmetry of the approximate custodial $SO(4)$ symmetry. In section V B we show that all these exact subsymmetries of the approximate custodial $SO(4)$ symmetry also are symmetries of the terms $\widehat{C}^2$, and hence there is no symmetry that can be imposed on the general C invariant NHDM, which expels the terms of the type $\widehat{C}^2$ only.

This paper is organized as follows: In section II we revise the custodial symmetry of the standard model (SM). In section III we reinvestigate the symmetry properties of the operators which are the building blocks of the NHDM potential, in terms of bidoublets—that is the Higgs fields represented by 4×4 complex matrices. Moreover we introduce some terminology, and quote some results from [22]. The subject of section IV is to what extent the $O(4)$-symmetry containing the custodial $SO(4)$-symmetry is a symmetry of the different parts of the Higgs and gauge field Lagrangian. Finally, in section V we search for symmetries beyond the global $SU(2)_L \times U(1)_Y$ for the kinetic Higgs terms, by investigating the adjoint action of $O(4)$ on the Lie algebra $u(2)$, or more concretely, the well-defined $O(4)$ transformations of the gauge bosons, see eq. (30) below and the discussion thereof. Some more mathematical discussions are delegated to the appendices.

II. THE APPROXIMATE $SO(4)$ SYMMETRY OF THE SM HIGGS LAGRANGIAN

Let the complex SM Higgs doublet be written

$$\Phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix}. \tag{2}$$

Then the global $SO(4) = (SU(2)_L \times SU(2)_R)/\mathbb{Z}_2$ custodial symmetry [21, 24] of the SM Higgs Lagrangian (in the limit $g' \to 0$) can be made manifest by rewriting the Higgs doublet as a matrix (bidoublet)

$$\check{\Phi} = \begin{pmatrix} \phi^{0*} & \phi^+ \\ -\phi^{+*} & \phi^0 \end{pmatrix} = \begin{pmatrix} \eta - i\phi_3 & \phi_1 + i\phi_2 \\ -\phi_1 + i\phi_2 & \eta + i\phi_3 \end{pmatrix}. \tag{3}$$

Here the Higgs potential will be a function of $\mathrm{Tr}[\check{\Phi}^\dagger\check{\Phi}]$, and is hence invariant under the global transformation

$$\check{\Phi} \to U_L \check{\Phi} U_R, \tag{4}$$

where U_L and U_R are $SU(2)$ matrices. The matrix U_L represents the usual gauged $SU(2)_L$ invariance. On the other hand, U_R represents an ordinary global transformation (where the gauge fields, when $g' \to 0$, does not transform in parallel with the Higgs doublet).

The SM scalar Lagrangian density can then be written [20]

$$\mathcal{L} = \frac{1}{2}\left(\mathrm{Tr}[(D_\mu\check{\Phi})^\dagger D^\mu\check{\Phi}] + \mu^2\mathrm{Tr}[\check{\Phi}^\dagger\check{\Phi}] - \lambda\mathrm{Tr}[\check{\Phi}^\dagger\check{\Phi}\check{\Phi}^\dagger\check{\Phi}]\right), \tag{5}$$

where the covariant derivative in the present notation is

$$D_\mu\check{\Phi} = \partial_\mu\check{\Phi} + \frac{1}{2}igW_{i\mu}\sigma_i\check{\Phi} - \frac{1}{2}ig'B_\mu\check{\Phi}\sigma_3, \tag{6}$$

where σ_i are the Pauli matrices. We see that the last term breaks the $SU(2)_R$ symmetry because of the factor σ_3. However, in the limit $g' \to 0$ the whole SM scalar Lagrangian has the full $SU(2)_L \times SU(2)_R$ symmetry, when the W fields transform (as usual) as a triplet under (the gauged) $SU(2)_L$ and as a singlet under the global $SU(2)_R$ symmetry. Some authors refer to $SU(2)_R$ as the custodial symmetry [25], others use the term of the symmetry $SU(2)_{L+R}$ which leaves the SM vacuum invariant [24]. In this article we denote the $SO(4) \cong (SU(2)_L \times SU(2)_R)/\mathbb{Z}_2$ symmetry as the custodial $SO(4)$ symmetry, as in [26].

III. THE OPERATORS $\widehat{B}$ AND $\widehat{C}$ OF THE NHDM IN THE BIDOUBLET FORMULATION $\check{\Phi}$

Consider the bilinear (i.e. linear in both Φ_m and Φ_n), hermitian operators $\widehat{B}$ and $\widehat{C}$ introduced in eq. (2.3) of [22], defined by[1]

$$\begin{aligned} \widehat{B}_{mn} = \widehat{B}(\Phi_m, \Phi_n) &= \tfrac{1}{2}(\Phi_m^\dagger \Phi_n + \Phi_n^\dagger \Phi_m) \\ \widehat{C}_{mn} = \widehat{C}(\Phi_m, \Phi_n) &= -\tfrac{i}{2}(\Phi_m^\dagger \Phi_n - \Phi_n^\dagger \Phi_m), \end{aligned} \tag{7}$$

where Φ_i refers to a doublet of the form (2), where every scalar field has an additional index i,

$$\Phi_i = \begin{pmatrix} \phi_i^+ \\ \phi_i^0 \end{pmatrix}, \quad i = 1, \ldots, N. \tag{8}$$

In (7) we let $1 \leq m \leq n \leq N$. The most general potential $V(\Phi_1, \ldots, \Phi_N)$ of the NHDM can then be built up by products and sums of the operators in (7). The most general C invariant NHDM potential can then be written

$$\begin{aligned} V_C(\Phi_1, \ldots, \Phi_N) = \mu_{mn} \widehat{B}_{mn} &+ \lambda^{(2)}_{mn,m'n'} \widehat{B}_{mn} \widehat{B}_{m'n'} \\ &+ \lambda^{(3)}_{mn,m'n'} \widehat{C}_{mn} \widehat{C}_{m'n'}, \end{aligned} \tag{9}$$

with an implicit sum over repeated indices, where $1 \leq m \leq n \leq N$ and $1 \leq m' \leq n' \leq N$ for terms containing $\widehat{B}$, and $1 \leq m < n \leq N$ and $1 \leq m' < n' \leq N$ for terms containing $\widehat{C}$. We also demand that the two pairs of indices $(mn)(m'n')$ are "lexicographically" ordered to avoid double counting. This means e.g. that indices $(12)(13)$ are included, while indices $(13)(12)$ are excluded in the sums of eq. (9). See sec. 2 in [22] for a more compact indexing. Superscripts in the parameters λ are chosen to coincide with the notation in [22].

The corresponding NHDM Lagrangian density is then given by

$$\mathcal{L}(x) = \sum_m [D^\mu \Phi_m(x)]^\dagger [D_\mu \Phi_m(x)] - V(\Phi_1, \Phi_2, \ldots, \Phi_N), \tag{10}$$

where the covariant derivative here is defined as

$$D^\mu = \partial^\mu + ig\frac{\sigma^i}{2} W_i^\mu + ig' Y B^\mu. \tag{11}$$

The term 'NHDM Lagrangian' is often used about the Lagrangian density (10). We will apply the term 'NHDM Lagrangian' as referring to the Lagrangian density (10) augmented by the gauge field Lagrangian given by eq. (33) below.

In the same manner as in the last section, we will now show that the operator $\widehat{C}$ does not share the $SO(4)$ symmetry held by the rest of the NHDM potential: Let $\check{\Phi}_i$ refer to a bidoublet of the form (3), where every scalar field has an additional index i. A simple calculation shows that

$$\widehat{B}(\Phi_m, \Phi_n) = \frac{1}{2} \mathrm{Tr}(\widehat{B}(\check{\Phi}_m, \check{\Phi}_n)), \tag{12}$$

while

$$\widehat{C}(\Phi_m, \Phi_n) = -\frac{1}{2} \mathrm{Tr}(\widehat{C}(\check{\Phi}_m, \check{\Phi}_n)\sigma_3) = -\frac{1}{2} \mathrm{Tr}(\sigma_3 \widehat{C}(\check{\Phi}_m, \check{\Phi}_n)). \tag{13}$$

---

[1] The operator $\widehat{A}_m$ defined in [22] equals $\widehat{B}_{mm}$ (no sum over m).



Figure 1: Diagram showing the overlap between relevant symmetry groups. The $SO(4)$ symmetry group is the custodial symmetry, $U(2) \cong SO(4) \cap Sp(2,\mathbb{R}) \cong SU(2)_L \times U(1)_Y$ is the global symmetry of the SM, $Sp(2,\mathbb{R})$ is the symmetry group of operators of type $\widehat{C}$, while $P(2,\mathbb{R})$ is the symmetry group of operators of type $\widehat{C}^2$. Finally, $O(4)$ is the symmetry group of operators of type $\widehat{B}$. Charge conjugation C will be an element of $P(2,\mathbb{R})^- \cap SO(4)$ where $P(2,\mathbb{R})^- = P(2,\mathbb{R}) - Sp(2,\mathbb{R})$, and the same is true for the matrix $\mathcal{J}$ of eq. (17) (with $k = 2$).

The latter confirms that the operator $\widehat{C}$, in contrast to $\widehat{B}$, does not have the $SU(2)_R$ [and hence neither the $SO(4)$-] symmetry since the presence of the factor σ_3 hinders us from utilizing the cyclic property of the trace. Furthermore, the identity

$$\begin{aligned}\widehat{C}(\Phi_m,\Phi_n)\widehat{C}(\Phi_{m'},\Phi_{n'}) &= -\frac{1}{2}\mathrm{Tr}(\widehat{C}(\check{\Phi}_m,\check{\Phi}_n)\sigma_3\widehat{C}(\check{\Phi}_{m'},\check{\Phi}_{n'})\sigma_3) \\ &= -\frac{1}{2}\mathrm{Tr}(\sigma_3\widehat{C}(\check{\Phi}_m,\check{\Phi}_n)\sigma_3\widehat{C}(\check{\Phi}_{m'},\check{\Phi}_{n'})),\end{aligned} \tag{14}$$

infers that operators of the type $\widehat{C}^2$, in the same manner as for $\widehat{C}$ above, do not share the $SO(4)$ symmetry of a NHDM potential built up by the operators of the types $\widehat{B}$ and $\widehat{B}^2$ only. This is also shown in section 2.2 of [22], but in a different manner: Here we showed that the symmetry group of $\widehat{C}$ is $Sp(2,\mathbb{R})$, while the symmetry group of $\widehat{C}^2$ is $P(2,\mathbb{R})$, with

$$P(2,\mathbb{R}) = Sp(2,\mathbb{R}) \cup P(2,\mathbb{R})^-. \tag{15}$$

Here the real symplectic group $Sp(2,\mathbb{R})$ is defined by

$$Sp(k,\mathbb{R}) = \{S \in GL_{2k}(\mathbb{R}) | S^T \mathcal{J} S = \mathcal{J}\}, \tag{16}$$

with

$$\mathcal{J} = \begin{pmatrix} 0_k & I_k \\ -I_k & 0_k \end{pmatrix}, \tag{17}$$

where 0_k and I_k is the $k \times k$ zero matrix and the $k \times k$ identity matrix, respectively. The component $P(2,\mathbb{R})^-$ of the Lie group $P(2,\mathbb{R})$, defined by

$$P(k,\mathbb{R})^- = \{S \in GL_{2k}(\mathbb{R}) | S^T \mathcal{J} S = -\mathcal{J}\}, \tag{18}$$

consists of matrices with determinant

$$\det(P(k,\mathbb{R})^-) = (-1)^k. \tag{19}$$

In the relevant case $k = 2$, P^- evidently consists of 4×4 matrices with determinant 1. The same is true for $Sp(2,\mathbb{R})$.

The custodial $SO(4)$ symmetry however, is not a subset of $P(2,\mathbb{R})$, and is hence not a symmetry of the operators of the type $\widehat{C}^2$ (nor $\widehat{C}$). See Fig. 1 for a diagram showing the intersections of the most important symmetry groups in this article.

IV. SYMMETRIES OF THE KINETIC TERMS

We now turn to the (global) symmetries of the kinetic terms of the NHDM Lagrangian,

$$K = \sum_{n=1}^{N} \left[(\partial^\mu + G^\mu)\,\Phi_n(x)\right]^\dagger \left[(\partial_\mu + G_\mu)\,\Phi_n(x)\right], \tag{20}$$

with

$$G^\mu = ig\frac{\sigma^i}{2}W_i^\mu + ig'YB^\mu. \tag{21}$$

We will let K_i denote the terms of the i'th order in the gauge fields.

A. Possible extensions of the global $SU(2)_L \times U(1)_Y$ symmetry

Now we will start investigating the possibility of having discrete or continuous symmetries beyond the global $SU(2)_L \times U(1)_Y \cong U(2)$ gauge symmetry[2], in the scalar sector. We do this by writing the NHDM Lagrangian in real form, i.e. we write the complex Higgs doublets as real quadruplets. In [22] we demonstrated that the kinetic terms of the NHDM Lagrangian has a symmetry group containing $SU(2) \times U(N)$ (enhanced to $SU(2) \times Sp(N)$ in the limit $g' \to 0$), when we allowed the different Higgs-fields to transform into each other. In the analysis below, we will not consider the possibility of different Higgs-doublets transforming into each other, since these transformations will not be symmetries of the general C invariant NHDM potential, see appendix A.

Consider the kinetic terms not involving gauge fields, K_0, of the Higgs Lagrangian,

$$K_0 = \sum_{n=1}^{N} \partial^\mu \Phi_n^T \partial_\mu \Phi_n, \tag{22}$$

where the complex Higgs doublet $(\Phi_n)_2 = \Psi_n + i\Theta_n$ is written in real form, as a quadruplet from now on denoted

$$\Phi_n = \begin{pmatrix} \Psi_n \\ \Theta_n, \end{pmatrix}. \tag{23}$$

We see that we can assign the terms K_0 a $O(4)$ symmetry by

$$\Phi_n \to O\Phi_n, \tag{24}$$

with $O \in O(4)$ and Φ_m given by eq. (23). The terms K_0 are invariant under the transformation (24) since $O^T O = I$ for $O \in O(4)$. The global symmetry $SU(2)_L \times U(1)_Y \cong U(2)$ will then be embedded into $SO(4) \subset O(4)$ by the map

$$\rho(X) = \begin{pmatrix} \mathrm{Re}(X) & -\mathrm{Im}(X) \\ \mathrm{Im}(X) & \mathrm{Re}(X) \end{pmatrix}, \tag{25}$$

cf. appendix B of [22]. Then the image of $U(2)$ under ρ,

$$\rho[U(2)] \subset SO(4), \tag{26}$$

is the global symmetry of the SM, when the fields are written as real quadruplets. The map ρ is an isomorphism onto its image $\rho[U(2)]$, and hence $\rho[U(2)] \cong U(2)$ (i.e. $\rho[U(2)]$ is a way to write $U(2)$ in real form).

We can then, as we showed in [22], write

$$\begin{aligned} K_1 &= \sum_{n=1}^{N} \partial^\mu (\Phi_n)_2^\dagger G_\mu (\Phi_n)_2 + (\Phi_n)_2^\dagger G^{\mu\dagger} \partial_\mu (\Phi_n)_2 \\ &= \sum_{n=1}^{N} \partial^\mu \Phi_n^T \mathcal{T}_\mu \Phi_n + \Phi_n^T(-\mathcal{T}^\mu)\partial_\mu \Phi_n \end{aligned} \tag{27}$$

where the subscript 2 in $(\Phi_n)_2$ indicates this is the usual complex Higgs doublet, while Φ_n is the four-dimensional real vector (23). Furthermore, the 4×4 matrix $\mathcal{T}^\mu$,

$$\mathcal{T}^\mu = \rho(G^\mu), \tag{28}$$

---

[2] More precisely we have $SO(3) \times U(1) \cong (SU(2)_L \times U(1)_Y)/\mathbb{Z}_2 \cong U(2)$, where the divisor $\mathbb{Z}_2$ is necessary since multiplication by $-I$ can be expressed both by $SU(2)$ and $U(1)$. We can hence identify elements $(-U) \times U' \sim U \times (-U')$, where $U \in SU(2)_L$ and $U' \in U(1)_Y$. Only multiplication by $-I \in U(2)$ can be expressed both by $U(1)$ and $SU(2)$: Assume $U = \exp(i\alpha)I \in SU(2)$. Then $1 = \det(U) = \exp(2\alpha i)$, and hence $\alpha = \pi$, i.e. $\exp(i\alpha) = -1$. The direct product $SU(2)_L \times U(1)_Y$ is therefore in reality a double cover of $U(2)$.

is given by [22]

$$\mathcal{T}^\mu = \begin{pmatrix} gW_I^\mu & -gW_R^\mu - g'YB^\mu I_2 \\ gW_R^\mu + g'YB^\mu I_2 & gW_I^\mu \end{pmatrix}, \tag{29}$$

with $W_R^\mu = \sum_{i=1,3} W_i^\mu \frac{1}{2}\sigma^i$ where the sum involves the real Pauli matrices σ^1 and σ^3. Moreover, $W_I^\mu = iW_2^\mu \frac{1}{2}\sigma^2$, where σ^2 is the imaginary Pauli matrix. The matrix I_2 is the 2×2 identity matrix.

We then see that the kinetic terms K_1 *apparently* are invariant under the $O(4)$ transformation (24) if we let the matrix $\mathcal{T}^\mu$ transform

$$\mathcal{T}^\mu \to O\mathcal{T}^\mu O^T \tag{30}$$

simultaneously with (24). As we will see, can the transformation (30) be described as the adjoint action of $O(4)$ on the Lie algebra $u(2)$ (the latter is represented by the matrix $\mathcal{T}^\mu$). The SM gauge bosons transform as usual under the global $SU(2)_L \times U(1)_Y \cong U(2) \cong \rho[U(2)] \subset SO(4)$, since for $u \in U(2)$

$$uG^\mu u^\dagger = \rho(u)\rho(G^\mu)\rho(u^\dagger) = \rho(u)\mathcal{T}^\mu\rho(u)^T, \tag{31}$$

where $uG^\mu u^\dagger$ is the way the gauge fields transform (globally) in the SM (they transform under the adjoint representation of $SU(2)_L \times U(1)_Y$) [30]. We will in sec. V B 1 note that in the case $g' \to 0$, the combined transformations (24) and (30) give us the custodial $SO(4)$ symmetry.

The only problem is that the transformation (30) might not be well-defined for choices of O beyond the global gauge group $U(2)$. The transformation is only well-defined when it induces consistent transformations of each of the fields; i.e. it is well-defined when it makes each of the fields transform in the same manner everywhere in the matrix $\mathcal{T}$. For instance, we cannot accept that B^μ transforms as $B^\mu \to B^\mu$ in $\mathcal{T}^\mu_{14}$, while it transforms as $B^\mu \to -B^\mu$ in $\mathcal{T}^\mu_{41}$. In section V we will investigate what kinds of transformations O make (30) well-defined.

Third, we consider the kinetic terms quadratic in the gauge fields,

$$\begin{aligned} K_2 &= \sum_{n=1}^N (\Phi_n)_2^\dagger G^{\mu\dagger} G_\mu (\Phi_n)_2 \\ &= -\sum_{n=1}^N \Phi_n^T \mathcal{T}^2 \Phi_n, \end{aligned} \tag{32}$$

which obviously are invariant under the $O(4)$ symmetry given by eqs. (24) and (30), when the latter transformation is well-defined.

B. The gauge field Lagrangian

In this section we want to show that transformation (30) also is a symmetry of the kinetic part of the gauge field Lagrangian. The reason for this is to ensure that symmetries contained in $O(4)$ are not violated by higher order diagrams involving diagrams generated by the (kinetic part of the) gauge field Lagrangian.

Consider the kinetic terms of the gauge field Lagrangian, formulated as the trace of the commutator of two covariant derivatives [as given in eq. (11)],

$$\mathcal{L}_{GB} = -\frac{1}{2}\mathrm{Tr}\left(\left(\frac{i}{g}[D_\mu, D_\nu]\right)^2\right)\Big|_{g'Y\to g}. \tag{33}$$

A general relation is

$$\mathrm{Tr}(\rho(X)) = 2\mathrm{Re}(\mathrm{Tr}(X)) = 2\mathrm{Tr}(X), \tag{34}$$

where the last equality is valid when the trace is real, as it is in eq. (33).

Then

$$\begin{aligned} \mathcal{L}_{GB} &= -\frac{1}{4}\mathrm{Tr}\left(\rho\left(\left(\frac{i}{g}[D_\mu, D_\nu]\right)^2\right)\right)\Big|_{g'Y\to g} \\ &= \frac{1}{4g^2}\mathrm{Tr}\left([\rho(D_\mu), \rho(D_\nu)]^2\right)\Big|_{g'Y\to g} \\ &= \frac{1}{4g^2}\mathrm{Tr}\left([\mathcal{T}_\mu, \mathcal{T}_\nu]^2\right)\Big|_{g'Y\to g} \end{aligned} \tag{35}$$

since ρ preserves matrix multiplication and addition, and since the partial derivatives commute. Eq. (35) is invariant under the transformation (30), by the cyclic property of the trace.

C. Yukawa couplings

The most general Yukawa couplings of the quark sector are of the form [17]

$$\mathcal{L}^{QH} = -\bar{Q}_{3,L}\left(\Gamma_{3\times 3,j}\Phi_{2,j}n_{3,R} + \Delta_{3\times 3,j}\tilde{\Phi}_{2,j}p_{3,R}\right) + \text{h.c.} \tag{36}$$

where summation over $j = 1, 2, \ldots, N$ is implicit. The symbol $\bar{Q}_L$ denotes a 1×3 vector consisting of the three left-handed quark doublets,

$$\bar{Q}_{3,L} = \left[(u_L^\dagger, d_L^\dagger)\ \ (c_L^\dagger, s_L^\dagger)\ \ (t_L^\dagger, b_L^\dagger)\right]\gamma^0, \tag{37}$$

where n_R and p_R denote (following the notation of [17]) 3×1 vectors consisting of the right-handed quark fields,

$$n_{3,R} = \begin{bmatrix} d_R \\ s_R \\ b_R \end{bmatrix}, \quad p_{3,R} = \begin{bmatrix} u_R \\ c_R \\ t_R \end{bmatrix}, \tag{38}$$

and where each of these six quark fields is a four-component Dirac spinor. Moreover, the complex doublet $\tilde{\Phi}_{2,j}$ is defined by

$$\tilde{\Phi}_{2,j} = -i(\Phi_{2,j}^\dagger \sigma_2)^T. \tag{39}$$

The authors of [27–29] has in the context of the 2HDM shown that the imposition of the custodial symmetry on the Yukawa couplings yields constraints on the coupling matrices Π_j and Δ_j, depending on the choice of the scalar field basis. After a possible change of scalar field basis[3], this corresponds to the case where no $O(4)$-violating operators of the type $\widehat{C}_{12}$ (or $\widehat{C}_{12}^2$, the latter equivalent to the constraint $\lambda_4 = \lambda_5$ in [29]) are present in the potential, and where [29]

$$\Pi_j = \Delta_j^\dagger, \tag{40}$$

for $j = 1, 2$. In the NHDM the custodial symmetry will infer restrictions similar to (40) on the Yukawa coupling matrices.

Imposing the custodial symmetry on the Yukawa coupling terms of the leptonic sector will lead to constraints analogous to (40) on the coupling matrices of the leptonic sector. Such restrictions will force the elements of the coupling matrices of the charged leptons and the neutrinos to be of the same order, making it hard to implement their great mass difference.

V. THE ADJOINT REPRESENTATION OF $O(4)$

To decide which transformations beyond $U(2)$ make the transformation (30) well-defined, we have to consider the adjoint representation of $O(4)$. We know the standard model gauge fields transform as the adjoint representation of $U(2)$, which is, written in real form, a sub-representation of the adjoint representation of $O(4)$. For a matrix $d \in O(4) - U(2)$ such that (30) is well-defined for $O = d$, the matrix d generates a discrete symmetry of the kinetic terms of the electroweak Lagrangian. Furthermore, if we could find such a $d \in O(4) - P(2,\mathbb{R})$ (where $P(2,\mathbb{R})$ is the symmetry group of the operators of the type $\widehat{C}^2$, defined in eq. (15), see also fig. 1), we could impose this discrete symmetry on the C invariant NHDM Lagrangian and hence avoid terms proportional to the $O(4)$-violating (but C invariant) operators $\widehat{C}^2$. Then we, because of the discrete symmetry, would have excluded these $O(4)$-violating terms both from the original potential and during renormalization.

---

[3] A change of scalar field basis is a transformation of the scalar fields, $\Phi_m \to \sum_n U_{mn}\Phi_n$, where $U \in U(N)$. Such a transformation changes the NHDM potential, but not the physics nor the symmetries thereof.

A. Some mathematical preliminaries

Generally, the adjoint representation of a (matrix) Lie group G is a homomorphism

$$Ad : G \to GL(\mathfrak{g}, \mathbb{R}), \tag{41}$$

where $\mathfrak{g}$ is the Lie algebra of G and

$$Ad(g)(X) = gXg^{-1}, \tag{42}$$

for $g \in G$ and $X \in \mathfrak{g}$. Ad is now a representation of G, acting on the vector space $\mathfrak{g}$. The linear transformation $Ad(g)$ on the Lie algebra $\mathfrak{g}$ is called the adjoint action of g on $\mathfrak{g}$. The set of all such linear transformations,

$$Ad[G] = \{Ad(g) \,:\, g \in G\}, \tag{43}$$

is called the adjoint action of (the Lie group) G on (its Lie algebra) $\mathfrak{g}$. Moreover, the set $GL(\mathfrak{g}, \mathbb{R})$ is the set of all linear, invertible transformations L, on the real vector space $\mathfrak{g}$. If the dimension of $\mathfrak{g}$ is n (i.e. $\mathfrak{g}$ has n basis vectors) $GL(\mathfrak{g}, \mathbb{R})$ consists of all real, invertible $n \times n$ matrices. In the case $G = SO(4)$, and hence $\mathfrak{g} = so(4)$ (which we will consider in the next section), the dimension of $\mathfrak{g}$ is $n = 6$, and therefore $GL(so(4), \mathbb{R})$ will consist of all real, invertible 6×6 matrices.

Denote the image $Ad[G]$ of the adjoint representation Ad_G, where $Ad_G \subset GL(\mathfrak{g}, \mathbb{R})$. When G is connected, the kernel of the adjoint representation is the center $Z(G)$ of G [31]

$$Z(G) = \{g \in G \,|\, \forall x \in G \,(xg = gx)\}. \tag{44}$$

The first isomorphism theorem of group theory then gives us

$$Ad_G \cong G/Z(G). \tag{45}$$

In the case of $SO(4)$, which is connected, $Z(SO(4)) = \{\pm I\}$. Hence

$$Ad_{SO(4)} = SO(4)/\{\pm I\}. \tag{46}$$

Now the group $SO(4)$ is not simple, in contrast to $SO(N)$ for $N = 3$ and for $N \geq 5$. Therefore $SO(4)$ can be, modulo its center $\{\pm I\}$, written as a direct product [35]

$$SO(4)/\{\pm I\} \cong SO(3) \times SO(3). \tag{47}$$

Hence, we have that

$$Ad_{SO(4)} \cong SO(3) \times SO(3), \tag{48}$$

which we also will see explicitly below.

B. The effect of the adjoint action

We will now explicitly consider the effect of the adjoint action of $O(4)$ (in fact on the Lie algebra $u(2)$, here parametrized by the gauge boson matrix $\mathcal{T}^\mu$), to see if it permits any symmetries beyond the (global) $U(2) = SU(2)_L \times U(1)_Y$ symmetry of the SM. We regard the real variant of $U(2)$; embedded in $SO(4)$ by the map ρ, and try to see if the adjoint action (30) can be well-defined for any matrices

$$O \in O(4) - U(2), \tag{49}$$

where $U(2)$ in reality is "shorthand" for $\rho[U(2)]$, the image of $U(2)$ under ρ. (We will do similar abbreviations many places in this article, as implied by the context.) The sets $U(2)$ and $\rho[U(2)]$ are the same Lie group, but the latter is expressed by real numbers. The matrix $\mathcal{T}$ only contains the gauge fields of the SM (W_1, W_2, W_3, B), that is, gauge fields corresponding to the Lie algebra $u(2) \subset so(4)$ (here $u(2)$ is written in real form, i.e. $u(2)$ is here shorthand for $\rho[u(2)]$). Hence the adjoint action (30) often will demand the extra gauge-fields X and Y of $o(4) - u(2)$, for all gauge fields to transform in consistent manners in $\mathcal{T}$, when the transformation (30) is carried out with an $O \in O(4) - U(2)$.

We will in the subsequent search for transformations $O \in O(4) - U(2)$ which do not demand the introduction of the extra gauge-fields of $o(4) - u(2)$ to make the adjoint action (30) consistent (i.e. well-defined).

A basis for the Lie algebra $so(4)$ is [22]

$$J_1 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad J_2 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad J_3 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix},$$

$$J_4 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix}, \quad J_5 = \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}, \quad J_6 = \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}. \tag{50}$$

Now regard the basis of the Lie algebra $so(4)$ given by

$$\begin{aligned} &\{X_i\} = \\ &\{\tfrac{1}{2}(J_2 + J_5), \tfrac{1}{2}(J_1 + J_4), \tfrac{1}{2}(J_6 - J_3), \tfrac{1}{2}(J_6 + J_3), \tfrac{1}{2}(J_2 - J_5), \tfrac{1}{2}(J_1 - J_4)\}, \end{aligned} \tag{51}$$

where $X_j = \rho\left((i/2)\sigma^j\right)$, $j = 1, \ldots, 3$, and $X_4 = \rho\left((i/2)I_2\right)$, i.e. the generators $X_1, \ldots, X_4$ are real forms of the generators of the $SU(2)_L \times U(1)_Y$ gauge group. Hence $X_1, \ldots X_4$ corresponds to the SM gauge bosons W_1, W_2, W_3, B, while X_5, X_6 would, if we regarded a full $SO(4)$ gauge symmetry, correspond to two non-SM gauge fields X and Y.[4] Then, since $\mathcal{T} = \rho(G)$ (here we suppress the Lorentz index μ),

$$\mathcal{T} = \sum_{j=1}^{4} w_j X_j, \tag{52}$$

where

$$w_j = g W_j \quad \text{for } j = 1, 2, 3, \quad \text{and} \quad w_4 = 2g' Y B, \tag{53}$$

and we also define

$$w_5 = g_X X, \quad w_6 = g_Y Y. \tag{54}$$

Since the Lie group $SO(4)$ is compact and connected, exponentiation of its Lie algebra generates the whole group: $\exp[so(4)] = SO(4)$. Moreover, we can write $SO(4)$ as a product of exponentiated generators ("one-parameter subgroups")

$$U = \{e^{t_1 X_1} e^{t_2 X_2} \cdots e^{t_6 X_6} \mid \vec{t} \in \mathbb{R}^6\} = SO(4). \tag{55}$$

Obviously $U \subset SO(4)$, since all the exponentials are elements of $SO(4)$. The equality $U = SO(4)$ will be demonstrated explicitly below, cf. the discussion following eq. (65).

We will now consider each exponentiation $e^{t_i X_i}$ of the generators X_i, and see which effect each of them has on the Lie algebra $so(4)$ under the adjoint action [the adjoint action will yield a linear transformation on so(4), according to (41)]:

Let $P_{ij}(\theta)$ denote the 6×6 matrix with elements

$$\begin{aligned} p_{ii} &= p_{jj} = \cos(\theta) \\ p_{ij} &= -p_{ji} = \sin(\theta) \\ p_{kk} &= 1 \quad k \neq i, j, \end{aligned} \tag{56}$$

---

[4] The Lie algebra $so(4) \cong su(2) \oplus su(2)$, and in our notation the matrices X_1, X_2, X_3 is a basis for one copy of $su(2)$ while X_4, X_5, X_6 is a basis for the other copy of $su(2)$. The matrices X_5, X_6 are chosen such that the usual commutator rules for the Lie algebra $su(2)$ is valid, for instance will $[X_5, X_6] = -X_4$, just as $[X_2, X_3] = -X_1$. Moreover, $[su(2) \oplus \{0\}, \{0\} \oplus su(2)] = [\sum_{j=1}^{3} t_j X_j, \sum_{j=4}^{6} t_j X_j] = 0$ for $t_1, \ldots, t_6 \in \mathbb{R}$, as the definition of a direct sum of Lie algebras demands.

with all other elements equaling zero. For instance, $P_{24}(\theta)$ is then given by

$$P_{24}(\theta) = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & \cos(\theta) & 0 & \sin(\theta) & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & -\sin(\theta) & 0 & \cos(\theta) & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}. \tag{57}$$

Let $so(4)$ be expressed by the basis $\{X_i\}$ of (51). Then, for example, the effect of $e^{t_1 X_1} \in SO(4)$ on $so(4)$ by the adjoint action is [see eqs. (30) and (42)],

$$e^{t_1 X_1} \sum_{i=1}^{6} w_i X_i e^{-t_1 X_1} = e^{t_1 \frac{1}{2}(J_2+J_5)} \sum_{i=1}^{6} w_i X_i e^{-t_1 \frac{1}{2}(J_2+J_5)} = E_1 \vec{w}\vec{X}, \tag{58}$$

where $\vec{w}\vec{X} = \sum_{i=1}^{6} w_i X_i$ is a general element in $so(4)$, and where the effect is summarized by the 6×6 matrix E_1. Hence $Ad(e^{t_1 X_1})$ maps $\vec{w}$ to $\vec{w'} = E_1 \vec{w}$.

Thus, the effect E_i of each $e^{t_i X_i}$ (no sum over i) on $so(4)$ by the adjoint action, can then be calculated by the formula $e^{t_i X_i} \vec{w}\vec{X} e^{-t_i X_i} = E_i \vec{w}$ to be

$$\begin{aligned} &E_1 = P_{23}(t_1),\ E_2 = P_{31}(t_2),\ E_3 = P_{12}(t_3) \\ &E_4 = P_{56}(t_4),\ E_5 = P_{64}(t_5),\ E_6 = P_{45}(t_6). \end{aligned} \tag{59}$$

Moreover, the effect E_r of the reflection

$$r = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} \tag{60}$$

is then

$$E_r = \begin{pmatrix} 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \end{pmatrix}. \tag{61}$$

The effect of the general element $u = e^{t_1 X_1} e^{t_2 X_2} \cdots e^{t_6 X_6}$ of $U \subseteq SO(4)$, U introduced in (55), is then given by

$$u \vec{w}\vec{X} u^{-1} = \vec{w'}\vec{X} \Leftrightarrow \vec{w'} = E\vec{w} \tag{62}$$

where

$$E = E_1 E_2 \cdots E_6. \tag{63}$$

Combining eqs. (59) and (63), we get

$$E = \begin{pmatrix} A(t_1, t_2, t_3) & 0_{3\times 3} \\ 0_{3\times 3} & A(t_4, t_5, t_6) \end{pmatrix}, \tag{64}$$

where

$$A(x, y, z) = \begin{pmatrix} c_y c_z & c_y s_z & -s_y \\ c_z s_x s_y - c_x s_z & c_x c_z + s_x s_y s_z & c_y s_x \\ c_x c_z s_y + s_x s_z & c_x s_y s_z - c_z s_x & c_x c_y \end{pmatrix}, \tag{65}$$

which is just the general element of $SO(3)$, written in the "xyz (pitch-roll-yaw) convention" [32]. Hence the effect E of the general element u of the set U in eq. (55) is just $SO(3) \times SO(3)$, since eq. (64) yields two independent copies of $SO(3)$. Since we already, in eq. (48), stated that $Ad_{SO(4)} = SO(3) \times SO(3)$, we know that the parametrization U of (55) covers the whole of $SO(4)$, at least modulo multiplication by $\pm I$, cf. eq. (47), where I denotes the identity. Moreover, since all exponentials $e^{t_j X_j}$ (no sum over j) consists of sines and cosines of angles $t_j/2$ we can express multiplication by minus the 4×4 identity, $-I_4$, in U;

$$e^{(t_j+2\pi)X_j} = -e^{t_j X_j} = -I_4 e^{t_j X_j}, \tag{66}$$

for any fixed $j \in \{1, \ldots, 6\}$. Hence $U = \pm U = SO(4)$, since all elements of all equivalence classes in $SO(4)/\{\pm I\}$ are elements of U, cf. eqs. (47) and (55).

We do not want to introduce gauge bosons beyond the SM, neither before nor after application of the adjoint action. Hence the parameters w_5, w_6 and w'_5, w'_6 must be zero (w_5, w'_5 corresponded to a non-SM gauge-boson X through the generator X_5, while w_6, w'_6 corresponded to a non-SM gauge boson Y through the generator X_6, see eq. (51) and the discussion after). Then we have to demand that some of the elements of E equal zero, namely

$$E_{5i} = E_{6i} = 0, \quad i = 1, \ldots, 4. \tag{67}$$

We want to find for which values of the angles $t_1, \ldots, t_6$ the effect of the adjoint action E does not force us to introduce non-SM gauge bosons X and Y, corresponding to $w_5 = g_X X$ and $w_6 = g_Y Y$. It is then sufficient to consider a matrix

$$\tilde{E} = E_5 E_6 = E(t_1 = \ldots = t_4 = 0), \tag{68}$$

for if

$$\vec{w'} = \tilde{E}\vec{w}, \tag{69}$$

with

$$w_5 = w_6 = w'_5 = w'_6 = 0, \tag{70}$$

then the matrix $E_1 \cdots E_4$,

$$E_1 \cdots E_4 = E(t_5 = t_6 = 0) = \begin{pmatrix} A(t_1, t_2, t_3) & 0_{3\times3} \\ 0_{3\times3} & A(t_4, 0, 0) \end{pmatrix}, \tag{71}$$

with

$$A(t_4, 0, 0) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{t_4} & s_{t_4} \\ 0 & -s_{t_4} & c_{t_4} \end{pmatrix}, \tag{72}$$

does not introduce non-SM gauge fields when applied on the vector $\vec{w'}$ of (69), that is,

$$\vec{w''} = E_1 \cdots E_4 \vec{w'}, \tag{73}$$

since $w''_5 = w''_6 = 0$.

Combining the expression for $\tilde{E}$ of eq. (68) and the relation $\vec{w'} = \tilde{E}\vec{w}$ for this case ($w_5 = w_6 = w'_5 = w'_6 = 0$) gives us the equation

$$\begin{pmatrix} w'_1 \\ w'_2 \\ w'_3 \\ w'_4 \\ 0 \\ 0 \end{pmatrix} = \vec{w'} = \tilde{E}\vec{w} = \begin{pmatrix} I_{3\times3} & & 0_{3\times3} & \\ & c_{t_5}c_{t_6} & c_{t_5}s_{t_6} & -s_{t_5} \\ 0_{3\times3} & -s_{t_6} & c_{t_6} & 0 \\ & c_{t_6}s_{t_5} & s_{t_5}s_{t_6} & c_{t_5} \end{pmatrix} \begin{pmatrix} w_1 \\ w_2 \\ w_3 \\ w_4 \\ 0 \\ 0 \end{pmatrix}, \tag{74}$$

We hence see that the only non-trivial constraints are $\tilde{E}_{54} = \tilde{E}_{64} = 0$.

The constraints $\tilde{E}_{54} = \tilde{E}_{64} = 0$ read

$$\begin{aligned} -\sin(t_6) &= 0, \\ \cos(t_6)\sin(t_5) &= 0, \end{aligned} \tag{75}$$

and hence,

$$\begin{aligned}\sin(t_5) &= 0,\\ \sin(t_6) &= 0.\end{aligned} \tag{76}$$

So we have

$$t_5, t_6 = n\pi, \quad n \in \mathbb{Z}, \tag{77}$$

not necessarily with the same n for both t_5 and t_6.

Now we want to check if the constraints (77) are consistent with any (discrete) symmetries $d \in SO(4) - P(2,\mathbb{R})$, i.e. symmetries beyond the symmetry group of the operators of the type $\widehat{C}^2$. The general element of $SO(4)$ can be written

$$O = e^{t_1 X_1} e^{t_2 X_2} e^{t_3 X_3} e^{t_4 X_4} e^{t_5 X_5} e^{t_6 X_6}, \tag{78}$$

but it is enough to consider matrices[5]

$$O = e^{t_5 X_5} e^{t_6 X_6} = \begin{pmatrix} c_5 c_6 & c_5 s_6 & s_5 s_6 & c_6 s_5 \\ -c_5 s_6 & c_5 c_6 & -c_6 s_5 & s_5 s_6 \\ -s_5 s_6 & c_6 s_5 & c_5 c_6 & -c_5 s_6 \\ -c_6 s_5 & -s_5 s_6 & c_5 s_6 & c_5 c_6 \end{pmatrix}, \tag{79}$$

where we have defined

$$c_i \equiv \cos(t_i/2), \quad s_i \equiv \sin(t_i/2). \tag{80}$$

We hence find that the matrix O of eq. (79) depends on angles $t_i/2$ for $i = 5, 6$. Then we have to regard the cases where

$$t_5, t_6 \in \{0, \pi, 2\pi, 3\pi\}, \tag{81}$$

cf. eq. (77). We then find that

$$\begin{gathered}\cos(t_5/2)\sin(t_6/2) = \pm 1 \quad \text{or} \quad \sin(t_5/2)\cos(t_6/2) = \pm 1\\ \Rightarrow O \in SO(4) \cap P(2,\mathbb{R})^-,\end{gathered} \tag{82}$$

while we otherwise (e.g. for $t_5 = t_6 = 0$) have

$$O \in SO(4) \cap Sp(2,\mathbb{R}) = U(2). \tag{83}$$

Eqs. (82) and (83) are found by comparing the form of O, given by eqs. (79) and (81), with respectively the forms (B3) and (B5) given in Appendix B. The set $P(2,\mathbb{R})^-$ in eq. (82) is a component of the symmetry group of the operators of the form $\widehat{C}^2$, see eq. (15). By considering the actual values of the angles t_5 and t_6 instead of half the angles, eqs. (82) and (83) can equivalently be written

$$\cos(t_5)\cos(t_6) = -1 \Rightarrow O \in SO(4) \cap P(2,\mathbb{R})^-, \tag{84}$$

$$\cos(t_5)\cos(t_6) = 1 \Rightarrow O \in SO(4) \cap Sp(2,\mathbb{R}) = U(2). \tag{85}$$

Eq. (84) shows that we get symmetries beyond the gauge group $U(2)$ if and only if the field B_μ changes sign,

$$B_\mu \to -B_\mu, \tag{86}$$

since the parameters w_4', w_4 corresponding to the transformation of the gauge field B_μ are given by $w_4' = E_{44} w_4$, see eq. (74), where $E_{44} = \cos(t_5)\cos(t_6)$ (64). Moreover, eq. (85) shows that B_μ transforms as

$$B_\mu \to B_\mu, \tag{87}$$

---

[5] Let $u \in U(2) \subset P(2,\mathbb{R})$ and $d \in SO(4) - P(2,\mathbb{R})$. Then $ud \notin P(2,\mathbb{R})$: Assume the opposite, $ud \in P(2,\mathbb{R})$. Then $u^{-1}(ud) = d \in P(2,\mathbb{R})$, a contradiction. Hence, if $d = e^{t_1 X_1} \cdots e^{t_6 X_6} \notin P(2,\mathbb{R})$, then $ud \notin P(2,\mathbb{R})$ for $u = (e^{t_1 X_1} \cdots e^{t_4 X_4})^{-1} \in U(2)$. Therefore $d' = ud \notin P(2,\mathbb{R})$, where $d' = e^{t_5 X_5} e^{t_6 X_6}$, and we only have to consider matrices of the same form as d'.

under (global) $U(2) \cong SU(2)_L \times U(1)_Y$ transformations, as it should according to the SM.

Furthermore, in appendix C we show that the symmetry group component the symmetries of (84) belong to, can be written

$$SO(4) \cap P(2,\mathbb{R})^- = U(2)S \tag{88}$$

for any

$$S \in SO(4) \cap P(2,\mathbb{R})^-. \tag{89}$$

This is especially true for $S = C \in SO(4) \cap P(2,\mathbb{R})^-$, where C is the charge conjugation operator. We implemented $C \in SO(4) \cap P(2,\mathbb{R})^-$ as

$$C = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} \tag{90}$$

in [22]. This corresponds to the transformation $C(\Phi) = \Phi^\dagger$ of a complex scalar doublet. If we want to include the possibility of a complex phase, i.e. $C(\Phi) = e^{i\alpha}\Phi^\dagger$, where Φ is a complex Higgs doublet, we still get $C \in SO(4) \cap P(2,\mathbb{R})^-$ (this can be shown by an argument similar to the one in footnote 5). Hence the arguments below are unaltered by a possible introduction of a phase $\alpha \neq 0$.

Now, since (by appendix C)

$$SO(4) \cap P(2,\mathbb{R})^- = U(2)C, \tag{91}$$

while

$$SO(4) \cap Sp(2,\mathbb{R}) = U(2) \tag{92}$$

we get that the Lie group G defined by

$$G = SO(4) \cap P(2,\mathbb{R}), \tag{93}$$

can be written

$$G = U(2) \cup U(2)C. \tag{94}$$

Since both $U(2)$ and $U(2)C$ are two sets of symmetries of the kinetic Higgs terms, the Lie group G is a symmetry group of the kinetic Higgs terms. There are by eqs. (84) and (85) no symmetries of the kinetic Higgs terms of the SM beyond the group G. This result is also derived in [33]. The group G is also the maximal symmetry group of a NHDM Lagrangian, provided that the NHDM potential is complicated enough not to allow any HF symmetries (beyond an overall $U(1)$ transformation). Hence G is the largest symmetry group of a NHDM Lagrangian with a sufficiently complicated potential, e.g. the most general C-invariant NHDM Lagrangian, see appendix A. If we only regard the kinetic terms of the NHDM Lagrangian, the maximal symmetry group will be $SU(N) \times G$, where $SU(N)$ are the Higgs family symmetries. The $U(1)_Y$ hypercharge symmetry is, as we have seen, contained in G.

By eq. (47) we can write $SO(4) \cong SU(2)_L \times SO(3)$ and by footnote 4 the three generators of $SU(2)_L$ commutes with the three generators of $SO(3)$ ($\cong SU(2)_R/\mathbb{Z}_2$). Charge conjugation C will, along with $U(1)_Y$, be a part of $SU(2)_R$. Hence C and $U(1)_Y$ will commute with $SU(2)_L$. We can therefore write $G = U(2) \cdot \{I, C\} = SU(2)_L \times (U(1)_Y \cdot \{I, C\})$, where $U(1)_Y \cdot \{I, C\} \cong O(2)$, since $U(1)$ may be mapped onto $SO(2)$ by the map ρ given in eq. (25), while C may be mapped to an arbitrary matrix with determinant -1. Now we get that $G \cong SU(2) \times O(2)$, but since both $SU(2)$ and $O(2)$ can express multiplication of the fields by $(-I)$, we divide by $\mathbb{Z}_2$ to avoid a double covering of G, where G is given in eqs. (93) and (94). Hence we have

$$G \cong SO(3) \times O(2). \tag{95}$$

Since $SO(2)$ is a normal subgroup of $O(2)$ [$gSO(2)g^{-1} \subseteq SO(2)$ for all $g \in O(2)$], and since $O(2) \cong U(1)_Y \cdot \{I, C\} \cong SO(2) \cdot \mathbb{Z}_2$, $O(2)$ can also be written as the semidirect product $SO(2) \rtimes \mathbb{Z}_2$. Since $CU(2)C \subseteq U(2)$, $U(2)$ will be a normal subgroup of G, and we can hence also write

$$G \cong U(2) \rtimes \mathbb{Z}_2. \tag{96}$$

Note that G cannot be written as the direct product of $U(2)$ and $\mathbb{Z}_2$. In fact,

$$G \ncong U(2) \times H, \tag{97}$$

for any group H, since the center of G is finite while the center of $U(2) \times H$ will be infinite, see Appendix D.

However, we find no well-defined $SO(4)$ transformation of the kinetic Higgs terms beyond the symmetry group $P(2,\mathbb{R})$ of the operator $\widehat{C}^2$. Hence, there is no discrete nor continuous symmetry we can impose on the NHDM Lagrangian to exclude the terms of the type $\widehat{C}^2$. (Again, assuming the potential is complicated enough to exclude the possibility of HF symmetries beyond an overall $U(1)$ transformation. This is the case for the most general C invariant NHDM potential, cf. appendix A.)

1. *The custodial $SO(4)$ symmetry*

On the other hand, if we also set $g' = 0$ (i.e. $w_4 = w_4' = 0$, since these are the parameters corresponding to the generator X_4 of the $U(1)_Y$ gauge group, see eq. (51) and the subsequent discussion), we see from eqs. (74) and (71) that all symmetries are well-defined, and hence the whole $SO(4)$ is a symmetry group of the SM Higgs Lagrangian. This is the custodial $SO(4)$ symmetry from section II, demonstrated in an alternative manner (for the kinetic terms). The symmetry equals the canonical implementation [26] of the custodial symmetry, since it is defined for the SM (N=1), by eq. (24).

C. The adjoint action of elements of $O(4)^-$

We also want to consider the adjoint action of elements of $O(4)^-$, the orthogonal matrices with determinant -1, and see if it can yield any (discrete or continuous) symmetries of the kinetic terms. The general effect of the adjoint action of an element of $O(4)^-$ can be written[6]

$$E^- = E_r \cdot E_1 E_2 \cdots E_6, \tag{98}$$

cf. eq. (63), where E_r is given in eq. (61), and we calculate E^- to be of the form

$$E^- = \begin{pmatrix} 0_3 & D(t_4,t_5,t_6) \\ C(t_1,t_2,t_3) & 0_3 \end{pmatrix}, \tag{99}$$

where 0_3 is the 3×3 zero matrix, and where the exact expressions of the 3×3 matrices C and D are irrelevant for the argument below.

In case E^- represented a well-defined adjoint action (i.e. the non-SM parameters equal zero: $w_5' = w_6' = w_5 = w_6 = 0$), E^- would have the effect

$$\begin{pmatrix} w_1' \\ w_2' \\ w_3' \\ w_4' \\ 0 \\ 0 \end{pmatrix} = E^- \begin{pmatrix} w_1 \\ w_2 \\ w_3 \\ w_4 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} E^-_{14} w_4 \\ E^-_{24} w_4 \\ E^-_{34} w_4 \\ \sum_{j=1}^3 E^-_{4j} w_j \\ \sum_{j=1}^3 E^-_{5j} w_j \\ \sum_{j=1}^3 E^-_{6j} w_j \end{pmatrix}. \tag{100}$$

But then all gauge fields $W_j, j = 1,2,3$ (the Lorentz index μ is suppressed) are transformed into some real number times the field B, since $w_j = gW_j$ for $j = 1,\ldots,3$ and $w_4 = 2g'YB$. But this transformation has no inverse transforming the B's (w_4) back into the different W_j's ($w_j, j = 1,2,3$), since it would have transformed different instances of the field B in a different manner. Hence no such well-defined E^-, as it appears in eq. (100), exists, because E^- should have an inverse (an adjoint action is a representation of a group, and all elements of a group has an inverse). Therefore $O(4)^-$ does not provide any new symmetries of the kinetic Higgs terms of the SM (or NHDM) Lagrangian.

---

[6] $O(n)^- = SO(n)R = RSO(n)$ for any $R \in O(n)^-$: We have $RSO(n) \subset O(n)^-$ since $\det(R)\det(S) = -1$ for $S \in SO(n)$, and $O(n)^- \subset RSO(n)$ since for $O \in O(n)^-$, $\det(R^T O) = 1$, hence $R^T O \in SO(n)$, and then $O = R(R^T O) \in RSO(n)$. The effect under the adjoint representation of an arbitrary element $O \in SO(4)$ can be calculated by $E_1 E_2 \cdots E_6$, cf. eqs. (62) and (63). Then the effect of an arbitrary element $RO \in O(4)^-$ can be calculated by $E_r E_1 E_2 \cdots E_6$, cf. (61).

The component $O(4)^-$ of $O(4)$ does not provide any new symmetries in the limit $g' \to 0$ (i.e. $w_4 = w_4' = 0$) either: In this case $W_j, j = 1, 2, 3$ would have to be mapped to 0, so this mapping could have no inverse, hence it cannot be an element of (the adjoint representation of) $O(4)$. This means the custodial $SO(4)$ symmetry cannot be extended by any (discrete or continuous) symmetry in $O(4)$. The result is also derived in [33]. It is also proven by one of the results in [34]. In this article all 13 possible accidental symmetry groups of a 2HDM potential, together with the kinetic Higgs terms in the limit $g' \to 0$, are determined. Here the maximal symmetry group, in case of a potential consisting of all possible $O(4)$ symmetric terms of the (in our terminology) type $\widehat{B}$ and $\widehat{B}^2$ (but transformed into a specific basis, see appendix A), is determined to be $SO(3) \times SU(2)_L \cong SO(4)$. I.e. the custodial $SO(4)$ symmetry is the maximal symmetry group of the kinetic Higgs terms in the limit $g' \to 0$ (when we are not regarding $SU(N)$ HF-symmetries).

VI. SUMMARY

By studying which $O(4)$ transformations $\mathcal{T}^\mu \to O^T \mathcal{T}^\mu O$ [cf. eq. (30)] of the gauge bosons are well-defined, we found the maximal set of symmetry transformations of the kinetic Higgs terms of the SM to constitute the Lie group $G = SO(4) \cap P(2, \mathbb{R}) \cong SO(3) \times O(2)$. The maximal symmetry group of the kinetic terms of the NHDM was then $SU(N) \times G$. The Lie group $P(2, \mathbb{R})$ was the symmetry group of operators of the form $\widehat{C}^2$, that is the $O(4)$ violating terms in the general C invariant NHDM potential. Hence, we could find no discrete nor continuous symmetry that is a symmetry of the kinetic terms and the quadratic (in the Higgs fields) terms of the type $\widehat{B}$, while not a symmetry of the terms of the type $\widehat{C}^2$. To show this, we also used the fact that the Higgs family symmetries of a potential containing all terms $\mu_{mn}\widehat{B}_{mn}$, $m \leq n$, also are symmetries of the terms of the type $\widehat{C}^2$, cf. appendix A. This means there is no symmetry we can impose on the general NHDM Lagrangian, to exclude the $O(4)$ violating terms from the NHDM potential, and to prevent them occurring as counterterms during the renormalization procedure, when calculating scattering processes.

As implied above, we found there are no symmetries of the kinetic Higgs terms in the negative determinant component $O(4)^-$ of $O(4)$. This was so, even in the limit $g' \to 0$. Hence, the custodial $SO(4)$ symmetry cannot be extended by elements of $O(4)^-$.

Acknowledgments

The author wants to thank Kåre Olaussen, Per Osland and David Fleming for helpful conversation.

Appendix A: Higgs family symmetries

In this appendix we will show that the $U(N)$ Higgs family (HF) symmetries[7] [23] of a potential containing all terms $\mu_{mn}\widehat{B}_{mn}$, $m \leq n$, also are Higgs family symmetries of the bilinears of the type $\widehat{C}_{mn}$. Hence there is no HF symmetry we can impose on the general C invariant NHDM potential, which exactly excludes the $O(4)$ violating terms, i.e. the terms of the form $\widehat{C}^2$.

Under a HF transformation the bilinear $\widehat{B}_{kk} \equiv \Phi_k^\dagger \Phi_k$ is transformed in the following manner

$$\Phi_k^\dagger \Phi_k \to \sum_{m,n=1}^{N} (U_{km}\Phi_m)^\dagger (U_{kn}\Phi_n) = \sum_{m,n=1}^{N} U_{km}^* U_{kn} \Phi_m^\dagger \Phi_n. \tag{A1}$$

If this $U(N)$ transformation (i.e. HF symmetry) keeps $\Phi_k^\dagger \Phi_k$ invariant, then (no sum over k)

$$\begin{aligned} U_{km}^* U_{kn} &= 1, \qquad \text{when } m = n = k, \\ U_{km}^* U_{kn} &= 0, \qquad \text{otherwise.} \end{aligned} \tag{A2}$$

---

[7] Here we also consider the $U(1)$ factor [of $U(N)$] as a HF symmetry, in addition to the $SU(N)$ HF symmetries. Overall multiplication by an $U(1)$ factor is in fact already taken care of by $U(1)_Y$, which we usually consider as a gauge group symmetry and not a HF symmetry.

This is so because the set $\{\Phi_m^\dagger \Phi_n\}_{m,n\leq N}$ (with N^2 elements) is linearly independent: The set $\{\Phi_n^\dagger, \Phi_m\}$ can be regarded classically as a set of $2N$ independent functions. Assume that

$$\sum_{i,j=1}^{N} a_{ij}\Phi_i^\dagger \Phi_j \equiv 0. \tag{A3}$$

If we then differentiate by $\partial^2/(\partial\Phi_m^\dagger \partial\Phi_n)$, for arbitrary m and n, we get that $a_{mn} = 0$, i.e.

$$a_{mn} = 0 \quad \text{for all} \quad m, n. \tag{A4}$$

This again means that the set $\{\Phi_m^\dagger \Phi_n\}_{m,n\leq N}$ is linearly independent.

Eq. (A2) then gives us that the most general HF transformation which keeps a term $\mu_{kk}\widehat{B}_{kk}$ (no sum over k) invariant, $1 \leq k \leq N$, is

$$\Phi_k \to \exp(i\alpha_k)\Phi_k. \tag{A5}$$

When we want to keep the term $\mu_{kk}\widehat{B}_{kk}$ invariant for all k, each bilinear has to transform as given in eq. (A5), with a possible different phase α_k for each k.

Now, if the potential we are regarding also contains all terms of the type $\mu_{mn}\widehat{B}_{mn}$ for $m < n$, in addition to the terms $\mu_{kk}\widehat{B}_{kk}$, there will be further restrictions on the allowed HF symmetries. The bilinear $\widehat{B}_{mn}$, $m < n$, now transforms as

$$\widehat{B}_{mn} \equiv \tfrac{1}{2}(\Phi_m^\dagger \Phi_n + \Phi_n^\dagger \Phi_m) \to \tfrac{1}{2}(U_{mm}^* \Phi_m^\dagger U_{nn}\Phi_n + U_{nn}^* \Phi_n^\dagger U_{mm}\Phi_m) \tag{A6}$$

(no sum over m, n) by eq. (A2) [valid for all k]. When $\widehat{B}_{mn}$, $m < n$, is kept invariant, that is,

$$U_{mm}^* \Phi_m^\dagger U_{nn}\Phi_n + U_{nn}^* \Phi_n^\dagger U_{mm}\Phi_m - \Phi_m^\dagger \Phi_n - \Phi_n^\dagger \Phi_m = 0, \tag{A7}$$

(no sum over m, n), we must have

$$U_{mm}^* U_{nn} = 1, \tag{A8}$$

since the two terms $\Phi_m^\dagger \Phi_n$ and $\Phi_n^\dagger \Phi_m$ are linearly independent. Eq. (A8) infers that the corresponding angles are identical,

$$\alpha_m = \alpha_n, \tag{A9}$$

where $U_{kk} = \exp(i\alpha_k)$. Hence the HF symmetries that keep all terms $\mu_{mn}\widehat{B}_{mn}$ simultaneously invariant for all m, n with $1 \leq m \leq n \leq N$, is an $U(1)$ transformation

$$\Phi_j \to \Phi_j \exp(i\alpha), \tag{A10}$$

for all j, $1 \leq j \leq N$. But this transformation also leave all terms $\widehat{C}_{mn}$ and $\widehat{C}_{mn}\widehat{C}_{m'n'}$ invariant, and hence the transformation cannot be applied to remove terms of the type $\widehat{C}^2$.

The result in (A10) and above is consistent with a result in [34]. Here the maximal symmetry group of the general C invariant 2HDM potential (expressed in the diagonally reduced basis) is found to be the group generated by CP (i.e. C) and $U(1)_Y$.

The general 2HDM potential of [34] can be expressed by the bilinear operators $\widehat{B}$ and $\widehat{C}$, in terms of the parameters given in [34] as

$$\begin{aligned} V = & -\mu_1^2\widehat{B}_{11} - \mu_2^2\widehat{B}_{22} - 2\mathrm{Re}(m_{12}^2)\widehat{B}_{12} + 2\mathrm{Im}(m_{12}^2)\widehat{C}_{12} + \lambda_1\widehat{B}_{11}^2 + \lambda_2\widehat{B}_{22}^2 + \lambda_3\widehat{B}_{11}\widehat{B}_{22} \\ & + [\lambda_4 + \mathrm{Re}(\lambda_5)]\,\widehat{B}_{12}^2 + 2\mathrm{Re}(\lambda_6)\widehat{B}_{11}\widehat{B}_{12} + 2\mathrm{Re}(\lambda_7)\widehat{B}_{22}\widehat{B}_{12} - 2\mathrm{Im}(\lambda_5)\widehat{B}_{12}\widehat{C}_{12} \\ & - 2\mathrm{Im}(\lambda_6)\widehat{B}_{11}\widehat{C}_{12} - 2\mathrm{Im}(\lambda_7)\widehat{B}_{22}\widehat{C}_{12} + [\lambda_4 - \mathrm{Re}(\lambda_5)]\,\widehat{C}_{12}^2. \end{aligned} \tag{A11}$$

All parameters in eq. (A11) are real (in some cases by taking the real or imaginary part of a complex parameter). In the diagonally reduced basis, we can furthermore set $\mathrm{Im}(\lambda_5) = 0$ and $\lambda_6 = \lambda_7$. Then the potential symmetric under

C and $U(1)_Y$ (in addition to $SU(2)_L$) corresponds to the constraints $\mathrm{Im}(m_{12}^2)=0$ and $\mathrm{Im}(\lambda_6)=\mathrm{Im}(\lambda_7)=0$, i.e. we are left with the potential

$$\begin{aligned} V = &-\mu_1^2\widehat{B}_{11} - \mu_2^2\widehat{B}_{22} - 2\mathrm{Re}(m_{12}^2)\widehat{B}_{12} + \lambda_1\widehat{B}_{11}^2 + \lambda_2\widehat{B}_{22}^2 + \lambda_3\widehat{B}_{11}\widehat{B}_{22} \\ &+ [\lambda_4 + \mathrm{Re}(\lambda_5)]\,\widehat{B}_{12}^2 + 2\mathrm{Re}(\lambda_6)(\widehat{B}_{11}\widehat{B}_{12} + \widehat{B}_{22}\widehat{B}_{12}) + [\lambda_4 - \mathrm{Re}(\lambda_5)]\,\widehat{C}_{12}^2, \end{aligned} \tag{A12}$$

which is the most general C invariant 2HDM potential, expressed in the diagonally reduced basis. We note that it, according to both [34] and eq. (A10), has no HF symmetries (beyond $U(1)_Y$, when we consider this as a HF symmetry).

In the case $[\lambda_4 - \mathrm{Re}(\lambda_5)] = 0$, we see that the symmetry is enhanced, since the potential then only contains $O(4)$ symmetric operators (of the type $\widehat{B}$). When we include the Kinetic Higgs terms in the limit $g' \to 0$, the maximal total symmetry becomes $SO(4)$, cf. sec. V C. We also noted in sec. V C that this is consistent with a result in [34], which determines this 2HDM to have the maximal symmetry group $SO(3)\times SU(2)_L \cong (SU(2)\times SU(2)_L)/\mathbb{Z}_2 \cong SO(4)$. Finally we note that the greatest symmetry group of any 2HDM potential, with kinetic terms in the limit $g'\to 0$, is in [34] calculated to be $SO(5)\times SU(2)_L$. This symmetry corresponds to a potential with the additional restrictions $\mu_1^2=\mu_2^2$, $m_{12}^2=0$, $\lambda_1=\lambda_2$, $\lambda_3=2\lambda_1$ and $\lambda_j=0$ for $j=4,5,6$ and 7. This is in agreement with [22], where we determined the symmetry group of the kinetic terms in the limit $g'\to 0$ to be $Sp(N)\times SU(2)_L$, where $Sp(N)$ is the quaternionic symplectic group. The agreement is proven by the isomorphism $SO(5)\cong Sp(2)/\mathbb{Z}_2$ (see [35] p. 430), where the $\mathbb{Z}_2$ factor reflect that we can express multiplication of the fields by $-I$ both in $SU(2)_L$ and $Sp(2)$ (since $SU(2)_R \subseteq Sp(N)$).

Appendix B: The form of some $O(2k)$ matrices

Let $S\in O(2k)$ (the case $k=2$ is the most interesting for this article), and let

$$\mathcal{J}_k = \mathcal{J} = \begin{pmatrix} 0_k & I_k \\ -I_k & 0_k \end{pmatrix}. \tag{B1}$$

Then the condition $S^T\mathcal{J}S=\mathcal{J}$ [i.e. $S\in Sp(k,\mathbb{R})$] can be written

$$\begin{aligned} \mathcal{J}S &= (S^T)^{-1}\mathcal{J}, \\ \mathcal{J}S &= S\mathcal{J}, \end{aligned} \tag{B2}$$

which, by the definition (B1) of $\mathcal{J}$, forces the solutions of eq. (B2) to be exactly the matrices in $O(2k)$ of the form

$$S = \begin{pmatrix} A & B \\ -B & A \end{pmatrix}, \tag{B3}$$

for arbitrary $k\times k$ matrices A, B.

Again, let $S\in O(2k)$. Then the equation the condition $S^T\mathcal{J}S=-\mathcal{J}$ [which is the definition of $S\in P(k,\mathbb{R})^-$], can be written

$$\mathcal{J}S = -S\mathcal{J}, \tag{B4}$$

which forces the matrices S to be exactly the ones in $O(2k)$ of the form

$$S = \begin{pmatrix} A & B \\ B & -A \end{pmatrix}, \tag{B5}$$

for arbitrary $k\times k$ matrices A, B.

Appendix C: The component $SO(4)\cap P(2,\mathbb{R})^-$

In this appendix we want to prove that

$$U(2)S = SO(4)\cap P(2,\mathbb{R})^-, \tag{C1}$$

for any $S \in SO(4) \cap P(2,\mathbb{R})^-$.

First, we claim the set $P(2,\mathbb{R})^-$ is given by

$$P(2,\mathbb{R})^- = Sp(2,\mathbb{R})\,S = S\,Sp(2,\mathbb{R}), \tag{C2}$$

for any $S \in P(2,\mathbb{R})^-$:

$\mathbf{P(2,\mathbb{R})^- \subseteq Sp(2,\mathbb{R})S}$: Let $S' \in P(2,\mathbb{R})^-$. Then $S'S \in Sp(2,\mathbb{R})$ since

$$(S'S)^T \mathcal{J}(S'S) = S^T(-\mathcal{J})S = \mathcal{J}. \tag{C3}$$

Then $S' = TS$ for $T = S'S^{-1} \in Sp(2,\mathbb{R})$, i.e. $S' \in Sp(2,\mathbb{R})S$. This is true since $S', S^{-1} \in P(2,\mathbb{R})^-$ infer $S'S^{-1} \in Sp(2,\mathbb{R})$ by (C3). ($S^{-1} \in P(2,\mathbb{R})^-$ since $S \in P(2,\mathbb{R})^-$.) Similarly with $S\,Sp(2,\mathbb{R})$.

$\mathbf{P(2,\mathbb{R})^- \supseteq Sp(2,\mathbb{R})S}$: On the other hand, if $T \in Sp(2,\mathbb{R})$, then

$$(TS)^T \mathcal{J}(TS) = S^T \mathcal{J} S = -\mathcal{J}, \tag{C4}$$

so then $TS \in P(2,\mathbb{R})^-$. Similarly, $ST \in P(2,\mathbb{R})^-$.

Now we can derive eq. (C1): Let $S \in P(2,\mathbb{R})^- \cap SO(4)$. Then

$$\begin{aligned} U(2)S = (SO(4) \cap Sp(2,\mathbb{R}))S &= (SO(4)S) \cap (Sp(2,\mathbb{R})S) \\ &= SO(4) \cap P(2,\mathbb{R})^-, \end{aligned} \tag{C5}$$

the last equality by eq. (C2).

Appendix D: The center of G

In this appendix we will show that the Lie group $G = P(2,\mathbb{R}) \cap SO(4)$ has a finite center $Z(G) = \{\pm I_4\} \cong \mathbb{Z}_2$. Hence G cannot be isomorphic to a group with a infinite center, especially not groups of the form $U(2) \times H$, H arbitrary.

First, G is a Lie group since the intersection of two (topologically) closed subsets of $GL(n,\mathbb{R})$ is a (topologically) closed subset of $GL(n,\mathbb{R})$, and hence a Lie group. The center $Z(G)$ of G consists of the elements in G which commute with all elements in G, cf. eq. (44). Let $Y \in Z(G) \subset G$. Then

$$Y^T \mathcal{J} Y = \pm\mathcal{J}, \tag{D1}$$

since $Y \in P(2,\mathbb{R})$. Moreover, since Y commutes with any element in G, $Y^T Y \mathcal{J} = \pm\mathcal{J}$, and hence

$$Y^T Y = \pm I. \tag{D2}$$

Since $Y \in G \subset SO(4)$ we get $I = \pm I$, and then $+$ is the right sign in eq. (D1),

$$Y^T \mathcal{J} Y = +\mathcal{J}, \tag{D3}$$

which again infer that $Y \in Sp(2,\mathbb{R})$.

This means

$$Y \in Sp(2,\mathbb{R}) \cap SO(4) = U(2), \tag{D4}$$

and thus Y is an element of $U(2)$ which commutes with all elements of $G \supset U(2)$, and hence we also have

$$Y \in Z(U(2)). \tag{D5}$$

Now $Z(U(2)) \cong U(1)$, or more concretely

$$Z(U(2)) = \{\lambda I_2 : \lambda \in \mathbb{C} \wedge |\lambda| = 1\}, \tag{D6}$$

which is a consequence of Schur's lemma. Expressed as a 4×4 real matrix by the map ρ, an element $Y \in Z(U(2))$ is of the form

$$Y = \begin{pmatrix} \cos\theta I_2 & -\sin\theta I_2 \\ \sin\theta I_2 & \cos\theta I_2 \end{pmatrix}, \tag{D7}$$

where $\cos\theta = \mathrm{Re}(\lambda)$ and $\sin\theta = \mathrm{Im}(\lambda)$.

Since Y also is an element in $Z(G)$, it also needs to commute with all elements of G, especially $C \in P(2,\mathbb{R})^- \cap SO(4) \subset G$. This means $YC = CY$, that is,

$$\begin{pmatrix} c_\theta I_2 & -s_\theta I_2 \\ s_\theta I_2 & c_\theta I_2 \end{pmatrix} \begin{pmatrix} I_2 & 0_2 \\ 0_2 & -I_2 \end{pmatrix} = \begin{pmatrix} I_2 & 0_2 \\ 0_2 & -I_2 \end{pmatrix} \begin{pmatrix} c_\theta I_2 & -s_\theta I_2 \\ s_\theta I_2 & c_\theta I_2 \end{pmatrix} \tag{D8}$$

which yields $s_\theta \equiv \sin\theta = 0$ and $c_\theta \equiv \cos\theta = \pm 1$, which again give us

$$Y = \begin{pmatrix} \pm I_2 & 0_2 \\ 0_2 & \pm I_2 \end{pmatrix} = \pm I_4. \tag{D9}$$

Hence we have derived the result $Z(G) \subset \{\pm I_4\}$, and since both $\pm I_4$ evidently commute with all of G, we get

$$Z(G) = \{\pm I_4\} \cong \mathbb{Z}_2, \tag{D10}$$

i.e. a finite center.

On the other hand,

$$Z(U(2) \times H) = Z(U(2)) \times Z(H) \cong U(1) \times Z(H), \tag{D11}$$

which is infinite, since $U(1)$ is infinite. Thus we have $G \ncong U(2) \times H$ for any group H.

[1] G. Aad *et al.* [ATLAS Collaboration], “Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC,” Phys. Lett. B [arXiv:1207.7214 [hep-ex]].

[2] S. Chatrchyan *et al.* [CMS Collaboration], “Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC,” Phys. Lett. B [arXiv:1207.7235 [hep-ex]].

[3] F. Englert and R. Brout, “Broken symmetry and the mass of gauge vector bosons”, Phys. Rev. Lett. **13** (1964) 321;

[4] P. W. Higgs, “Broken symmetries and the masses of gauge bosons”, Phys. Rev. Lett. **13** (1964) 508;

[5] G. S. Guralnik, C. R. Hagen and T. W. B. Kibble, “Global Conservation Laws and Massless Particles,” Phys. Rev. Lett. **13** (1964) 585.

[6] P. W. Higgs, “Spontaneous Symmetry Breakdown Without Massless Bosons”, Phys. Rev. **145** (1966) 1156

[7] V. Silveira and A. Zee, “Scalar Phantoms”, Phys. Lett. B **161** (1985) 136.

[8] D. E. Holz and A. Zee, “Collisional dark matter and scalar phantoms”, Phys. Lett. B **517** (2001) 239 [arXiv:hep-ph/0105284].

[9] E. Ma, “Verifiable radiative seesaw mechanism of neutrino mass and dark matter”, Phys. Rev. D **73** (2006) 077301 [arXiv:hep-ph/0601225].

[10] R. Barbieri, L. J. Hall and V. S. Rychkov, “Improved naturalness with a heavy Higgs: An alternative road to LHC physics”, Phys. Rev. D **74** (2006) 015007 [arXiv:hep-ph/0603188].

[11] B. Patt and F. Wilczek, “Higgs-field portal into hidden sectors”, arXiv:hep-ph/0605188.

[12] R. A. Porto and A. Zee, “The Private Higgs”, Phys. Lett. B **666** (2008) 491 [arXiv:0712.0448 [hep-ph]].

[13] B. Grzadkowski, O. M. Ogreid and P. Osland, “Natural Multi-Higgs Model with Dark Matter and CP Violation”, Phys. Rev. **D80** (2009) 055013 [arXiv:0904.2173 [hep-ph]].

[14] I. P. Ivanov and V. Keus, “Z_p scalar dark matter from multi-Higgs-doublet models,” arXiv:1203.3426 [hep-ph].

[15] T. D. Lee, “A Theory of Spontaneous T Violation,” Phys. Rev. **D8** (1973) 1226-1239.

[16] S. Weinberg, “Gauge Theory of CP Violation,” Phys. Rev. Lett. **37** (1976) 657.

[17] G. Branco, L. Lavoura and J. Silva, ”*CP* Violation”, Oxford (1999).

[18] C. C. Nishi, “CP violation conditions in N-Higgs-doublet potentials,” Phys. Rev. D **74** (2006) 036003 [Erratum-ibid. D **76** (2007) 119901] [hep-ph/0605153].

[19] G. C. Branco, P. M. Ferreira, L. Lavoura, M. N. Rebelo, M. Sher and J. P. Silva, “Theory and phenomenology of two-Higgs-doublet models,” Phys. Rept. **516**, 1 (2012) [arXiv:1106.0034 [hep-ph]].

[20] R. K. Ellis in M. Krämer and F. J. P. Soler "Large Hadron Collider Phenomenology" The Scottish Universities Summer School in Physics (2004).

[21] S. Willenbrock, “Symmetries of the standard model,” arXiv:hep-ph/0410370.

[22] K. Olaussen, P. Osland, M. Aa. Solberg, “Symmetry and Mass Degeneration in Multi-Higgs-Doublet Models,” JHEP **1107** (2011) 020. [arXiv:1007.1424 [hep-ph]].

[23] P. M. Ferreira and J. P. Silva, “Discrete and continuous symmetries in multi-Higgs-doublet models,” Phys. Rev. D **78** (2008) 116007 [arXiv:0809.2788 [hep-ph]].

[24] P. Sikivie, L. Susskind, M. B. Voloshin and V. I. Zakharov, “Isospin Breaking In Technicolor Models,” Nucl. Phys. B **173** (1980) 189.

[25] H. Georgi, “Effective field theory,” Ann. Rev. Nucl. Part. Sci. **43** (1993) 209.

[26] C. C. Nishi, “Custodial SO(4) symmetry and CP violation in N-Higgs-doublet potentials,” Phys. Rev. D **83** (2011) 095005 [arXiv:1103.0252 [hep-ph]].

[27] A. Pomarol and R. Vega, “Constraints on CP violation in the Higgs sector from the rho parameter,” Nucl. Phys. B **413** (1994) 3 [hep-ph/9305272].

[28] H. E. Haber and A. Pomarol, “Constraints from global symmetries on radiative corrections to the Higgs sector,” Phys. Lett. B **302** (1993) 435 [arXiv:hep-ph/9207267].

[29] H. E. Haber and D. O’Neil, “Basis-independent methods for the two-Higgs-doublet model III: The CP-conserving limit, custodial symmetry, and the oblique parameters S, T, U,” Phys. Rev. D **83** (2011) 055017 [arXiv:1011.6188 [hep-ph]].

[30] O. Nachtmann, “Elementary Particle Physics: Concepts And Phenomena,” Berlin, Germany: Springer (1990) 559 p

[31] W. Ziller "Lie Groups. Representation Theory and Symmetric Spaces" University of Pennsylvania (2010) [www.math.upenn.edu/ wziller/math650/LieGroupsReps.pdf]

[32] Weisstein, Eric W. "Euler Angles." From MathWorld–A Wolfram Web Resource. [http://mathworld.wolfram.com/EulerAngles.html]

[33] Chapter 4 in M. Aa. Solberg, Dark matter candidates and their indirect detection, PhD-thesis, NTNU, 2010. http://urn.kb.se/resolve?urn=urn:nbn:no:ntnu:diva-11162

[34] A. Pilaftsis, “On the Classification of Accidental Symmetries of the Two Higgs Doublet Model Potential,” Phys. Lett. B **706** (2012) 465 [arXiv:1109.3787 [hep-ph]].

[35] J. F. Cornwell, “Group Theory In Physics. Vol. 2,” London, Uk: Academic (1984) 589 P. (Techniques Of Physics, 7)